

RESEARCH NOTES

NESTS OF *HIBANA GRACILIS* ARE REUSED BY  
*PHIDIPPUS CLARUS* IN WETLANDS OF  
NORTHEASTERN KANSAS

Typical nests of wandering spiders consist of small, tightly packed hollow bundles. In these nests spiders molt, retire for part of the day and often lay their eggs (Jackson 1979; Foelix 1982). Nests may be constructed on the ground in natural shelters such as leaf litter or woody debris

or within a plant canopy among easily folded leaves or other structures (Johnson 1992).

In wetlands of northeastern Kansas in early May of 1993 and 1994, I found many mating pairs of *Hibana gracilis* (Hentz) (Araneae, Anyphaenidae) within nests made in the small up-



Figure 1.—Mating pair of *Hibana gracilis* (Araneae: Anyphaenidae) in the uppermost expanded leaves of the milkweed plant.

permost expanded leaves of common milkweed (*Asclepias syriaca*) and sullivant's milkweed (*A. sullivantii*) (Fig. 1). Because these milkweed species may be more common in annually burned wetlands (Johnson & Knapp, in press), I quantified the density and diversity of *H. gracilis* nests in three annually burned wetlands. Each wetland was approximately 350–500 m<sup>2</sup> in area and contained approximately 5.4 *A. syriaca* or *A. sullivantii* stems/m<sup>2</sup> (Johnson, in press). *Hibana gracilis* nests were found on 72% of milkweed stems in 40 plots, making nest density among milkweed in these wetlands 3.88 nests/m<sup>2</sup>. Nests were observed in this density on either *A. syriaca* or *A. sullivantii* depending on which species was more centrally located within wetlands. Otherwise, *H. gracilis* nests were found within basal leaf sheaths of prairie cordgrass (*Spartina pectinata*) in a density of 0.4 nests/m<sup>2</sup> ( $n = 30$  m<sup>2</sup> plots of *S. pectinata*).

By late May, *H. gracilis* had abandoned most nests; but the majority of these nests were reoccupied by *Phidippus clarus* Keyserling (Araneae, Salticidae). In the same three wetlands described above, 64% ( $n = 40$ ) of former *H. gracilis* nests in milkweed stems were occupied by a female or a pair of *P. clarus*. Furthermore, these nests did not appear to have been altered in any way from the original structures made by *H. gracilis*. By mid-June 1994, *P. clarus* had also abandoned these milkweed leaf nesting sites.

This pattern of nest reuse represents an interesting interaction between wandering arachnids and may be linked to 1) shortage of suitable nest-

ing sites, 2) reoccupation of optimal nest sites, or 3) silk conservation. Young milkweed plants may represent ideal nesting sites which are shared by sympatric arachnids with similar hunting methods and prey choices, but allochronic mating periods.

#### ACKNOWLEDGMENTS

I thank Dr. Bruce Cutler for his contributions in species identification and helpful advice. I also thank the Konza Prairie LTER Staff.

#### LITERATURE CITED

- Foelix, R. F. 1982. Biology of Spiders. Harvard University Press, Cambridge.
- Jackson, R. R. 1979. Nests of *Phidippus johnsoni* (Araneae, Salticidae): characteristics, patterns of occupation, and function. *J. Arachnol.*, 7:47–58.
- Johnson, S. R. 1993. Jumping spiders nest in red plastic area marker flags in prairies of northeastern Kansas. *Prairie Nat.*, 25:275.
- Johnson, S. R. In press. Wind-induced leaf binding by *Spartina pectinata* onto *Asclepias syriaca* in northeastern Kansas. *Prairie Nat.*
- Johnson, S. R. & A. K. Knapp. In press. The influence of fire on *Spartina pectinata* wetland communities in a northeastern Kansas tallgrass prairie. *Canadian J. Bot.*
- Stephen R. Johnson:** Division of Biology, Ackert Hall, Kansas State University, Manhattan, Kansas 66506 USA.

*Manuscript received 31 October 1994, revised 22 January 1995.*